

**Morbidity and mortality due to Nipah or Nipah-like virus encephalitis in
WHO South-East Asia Region, 2001-2018**

Country: Bangladesh

Month/Year	Location	No. cases	No. deaths	Case Fatality Rate
April, May 2001	Meherpur	13	9	69%
January 2003	Naogaon	12	8	67%
Jan 2004	Rajbari	31	23	74%
Apr 2004	Faridpur	36	27	75%
Jan- Mar 2005	Tangail	12	11	92%
Jan-Feb 2007	Thakurgaon	7	3	43%
Mar 2007	Kushtia	8	5	63%
Apr 2007	Pabna, Natore and Naogaon	3	1	33%
Feb 2008	Manikgonj	4	4	100%
Apr 2008	Rajbari	7	5	71%
Jan 2009	Gaibandha, Rangpur and Nilphamari	3	0	0%
	Rajbari	1	1	100%
Feb-Mar 2010	Faridpur	8	7	87.50%
	Faridpur, Rajbari, Gopalganj,	8	7	87.50%
	Kurigram,	1	1	100%
Jan-Feb 2011	Lalmohirhat, Dinajpur, Comilla Nilphamari, Faridpur, Rajbari	44	40	91%
Jan 2012	Joypurhat	12	10	83%
Jan- Apr 2013	Pabna, Natore, Naogaon, Gaibandha, Manikganj	24	21	88%
Jan-Feb 2014	13 districts	18	9	50%
Jan-Feb 2015	Nilphamari, Ponchoghor, Faridpur, Magura, Naugaon, Rajbari	9	6	67%

Country: India

Month/Year	Location	No. cases	No. deaths	Case Fatality Rate
Feb 2001	Siliguri	66	45	68%
Apr 2007	Nadia	5	5	100%
May* 2018	Kerala	14	12	86%

*As of 24 May 2018