

# Prevalence of hypertension among the Bangladeshi adult population: a meta-analysis

Moniruzzaman<sup>i</sup>, Abu Taleb<sup>i</sup>, Shahadur Rahman<sup>i</sup>, Amitava Acharyya<sup>i</sup>,  
Ferdous Ara Islam<sup>i</sup>, MSA Mansur Ahmed<sup>i</sup> and M Mostafa Zaman<sup>ii</sup>

## Abstract

Hypertension has become a significant problem in many developing countries undergoing epidemiological transition. A meta-analysis covering studies up to 1994 reported a prevalence of 11.3% in the adult population of Bangladesh. We conducted a meta-analysis to estimate the current prevalence of hypertension among the adult Bangladeshi population. We searched MEDLINE and included studies published in professional journals between 1995 and 2009 on the prevalence of hypertension among the adult Bangladeshi population. We included population-based studies that had a clear definition of hypertension, and were conducted in adults ( $\geq 15$  years). We located twelve articles, of which six articles were excluded from the current analysis as they were conducted in specific population subgroups. To obtain the prevalence of hypertension, we conducted a meta-analysis of these studies and recalculated their 95% confidence intervals, if required, to obtain a pooled estimate. Five of the six studies were from rural areas and were heterogeneous in terms of age groups studied and definition of hypertension used. The pooled estimate for the prevalence of hypertension in 6430 adults was 13.5% (12.7%–14.3%). Our meta-analysis clearly demonstrates the high burden of hypertension among the adult population of rural Bangladesh and underscores the importance of instituting screening for asymptomatic individuals, especially in primary health-care settings.

## Introduction

Hypertension is recognized as a major contributor to the disease burden globally. Hypertension and its complications account for an estimated 9.4 million deaths every year.<sup>1</sup> It has become a significant problem in many developing countries undergoing epidemiological transition.<sup>2</sup> The higher the blood pressure, the greater the chances of heart attack, heart failure, stroke and kidney disease.<sup>3</sup> The World Health Organization (WHO) attributes hypertension, or high blood pressure, to be the leading cause of cardiovascular mortality.

Bangladesh is passing through a phase of epidemiological transition from communicable diseases to noncommunicable disease (NCDs) and currently has a double burden of disease.<sup>4</sup> This means that the prevalence of hypertension is modest now but will show a rising trend.

There is a lack of representative data on the prevalence of hypertension in the Bangladeshi population. One meta-analysis conducted by Zaman and Rouf in 1999 on the prevalence of

<sup>i</sup> Bangladesh Institute of Health Sciences (BIHS), the sister organization of BIRDEM, 125/1 Darussalam, Mirpur, Dhaka-1216

<sup>ii</sup> World Health Organization (WHO), Bangladesh

hypertension in the Bangladeshi adult population included studies from 1976 to 1994, and estimated the prevalence to be 11.3%.<sup>5</sup> Between 1995 and 2009, some more studies with small sample sizes were conducted,<sup>6-16</sup> which showed varying prevalence rates. Individually, these studies cannot provide sufficient information about the prevalence in the country due to the non-representativeness of the study populations as well as the small sample sizes. In order to efficiently utilize the information available from these surveys, we decided to conduct another meta-analysis of all previous population-based studies in Bangladesh for a more accurate estimation of the prevalence of hypertension. We present the results of this exercise in this paper.

## Methods

For this meta-analysis, we identified articles published from 1995 to 2009 on the prevalence of hypertension among the Bangladeshi adult population by conducting a MEDLINE search. We included those population-based studies that had a clear diagnostic definition of hypertension and were conducted in the age group  $\geq 15$  years. We located 12 articles,<sup>6-16</sup> among which 5 were excluded from the current analysis because they did not fulfil our selection criteria. These were studies done in clinical settings,<sup>13</sup> on diabetic subjects,<sup>8</sup> pregnant women,<sup>14</sup> subjects with arsenicosis,<sup>15</sup> and among the elderly population ( $>60$  years).<sup>10</sup> If there was more than one publication from the same survey, we included only one of the publications and so we did not include the data of one study<sup>17</sup> (Table 1). We then calculated the prevalence of hypertension in these studies and the 95% confidence intervals (CI) for the individual studies using the data presented. Finally, we added the denominators (total number of subjects studied) and numerators (total number of subjects identified with hypertension) from each study to get the pooled prevalence estimate with its 95% CI. This meant that the weight given to each study was proportional to its sample size.

## Results

Table 1 summarizes the studies on the prevalence of hypertension in the Bangladeshi adult population from 1995 to 2009, and the calculated prevalence rates and their 95% confidence intervals. Except for one study, which was done in both urban and rural areas, the remaining five studies were done in rural Bangladesh and thus the results can be used as an estimate only for rural Bangladesh. Five studies included adults  $\geq 18$  years of age and the sixth one included persons  $\geq 15$  years. The prevalence of hypertension in these studies ranged from 7.8% to 18.6%.

All the studies did not use the definition of hypertension proposed by WHO in 1993, which defines hypertension as systolic and/or diastolic blood pressure ( $\geq 140/90$  mmHg) with or without a history of taking antihypertensive medication.

Table 1 provides the pooled estimate of the prevalence of hypertension as measured from the studies that met the eligibility criteria for this meta-analysis. From a total of 6430 adults included in these studies, we arrived at an estimate of 13.5% (with 95% CIs ranging from 12.7% to 14.2%).

## Discussion

Through this meta-analysis, we attempt to provide information on the prevalence of hypertension in the Bangladeshi adult population. For this purpose, we searched articles published in MEDLINE during the years 1995–2009 on the prevalence of hypertension in Bangladeshi adults.

Table 1. Articles on hypertension published from 1995 to 2009 identified by MEDLINE search

First author Year of publication (ref)	Place of study Year of study	Subjects	Sampling procedure	Response rate (%)	Number of readings taken	Criteria for diagnosis of hypertension	Age group (years)	Sample size (N)	Prevalence (%)	Meta-analysis		
										Age (years)	No. (N)	P (%)
Sayed A 1995 <sup>1</sup>	Dohar Not mentioned	Rural community of Dohar	Cluster sampling	70	Mean of three readings	SBP $\geq$ 140 mmHg DBP $\geq$ 90 mmHg	$\geq$ 15	1005	SBP 10.5 and DBP 9.0	1005	10.5	8.6-12.4
Sayed MA 1998 <sup>7</sup>	Bangladesh Not mentioned	NIDDM and IGT subjects	Random sampling	Not reported	Not reported	SBP $\geq$ 140 mmHg DBP $\geq$ 90 mmHg	30-60	693	SBP 23.2 and DBP 13.6	Not included		
Rahman M 1999 <sup>8</sup>	4 villages: Samta, Teekhalo, Rajerampur and Sonakhendi Not mentioned	Rural inhabitants	Multistage cluster sampling	100	Three readings; lowest was taken	SBP $\geq$ 140 mmHg DBP $\geq$ 90 mmHg combined with DBP $\geq$ 90 mmHg	$\geq$ 30	114	7.80	114	7.80	2.9-12.7
Zaman MM 2001 <sup>9</sup>	Rural area, Savar, Dhaka 1996	Rural population of Bangladesh	Simple random sampling	63	Mean of two readings	SBP $\geq$ 140 mmHg and/or DBP $\geq$ 90 mmHg or medication	$\geq$ 18	510	12.8	510	12.8%	9.9-15.7
Quasem I 2001 <sup>10</sup>	Mymensingh municipal and Muktadasa Thana, Mymensingh 1999-2000	Elderly population	Multistage cluster sampling	Not reported	Mean of two readings but three readings taken if > 10 mmHg variation found	WHO International Society for Hypertension criteria	> 60	480	Urban 7.5 and Rural 5.3	Not included		
Sayed MA 2002 <sup>11</sup>	Rural and urban community Not mentioned	Rural and urban community	Simple random sampling in rural and cluster sampling in urban	Rural 83.6 and urban 77.8	Mean of two readings	Not mentioned	> 20	2361	SBP 14.4 and DBP 9.1	2361	14.4	8.4-20.4
Zaman MM 2004 <sup>12</sup>	Rural community, Maitab, Chaudpur district 1999-2001	Clinic- based rural population	All	100	One reading	SBP $\geq$ 140 mmHg DBP > 90 mmHg	> 20	1271	17.80	Not included		
Sayed MA 2005 <sup>13</sup>	Randomly selected 10 villages of Mymensingh 2001-2002	Pregnant women	Not reported	Not reported	Not reported	WHO criteria was used	18-44	147	SBP 6.8 and DBP 5.4	Not included		
Chen Y 2006 <sup>14</sup>	Aratthazar 2000-2002	Rural Bangladeshis with exposure to arsenic	Not mentioned	97.50	Three readings; the lowest one was taken	SBP $\geq$ 140 mmHg or DBP $\geq$ 90 mmHg or antihypertensive drugs	$\geq$ 18	11116	13.30	Not included		
Zaman MM 2007 <sup>15</sup>	Eklbaur village, Chandpur district 2001	Rural Bangladeshis	Simple random sampling	64	One	Blood pressure $\geq$ 140/80 mmHg or antihypertensive drugs	$\geq$ 20, mean age 40	700, actual 447	18.60	440	18.6	15.0-22.2
Van Minh H 2009 <sup>16</sup>	Maitab, Mirsarai, Abhaynagar and WATCH area 2005	Rural Bangladeshis	Multistage cluster sampling	98	Mean of three readings	WHO criteria were used	25-64	2000	13.40	2000	13.4	11.9-14.9
<b>Pooled estimate of hypertension</b>										6430	13.5	12.7-14.2

NIDDM non-insulin dependent diabetes mellitus  
 IGT impaired glucose tolerance  
 SBP systolic blood pressure  
 DBP diastolic blood pressure

On analysis, we found a prevalence of hypertension of 13.5% in the adult population of Bangladesh, which was higher than the 11.3% prevalence reported by Zaman and Rouf in 1999.<sup>5</sup> The meta-analysis by Zaman and Rouf included studies conducted from 1976 to 1994 and the higher estimate reported in our study could represent a temporal trend. However, the previous meta-analysis included studies conducted in Dhaka only and mostly urban areas, where the prevalence of hypertension has been found to be higher, while our meta-analysis included studies conducted mainly among the rural community, which report a lower prevalence and would, therefore, probably have underestimated the increase in the prevalence of hypertension.

In the NCD risk factor survey conducted in Bangladesh in 2010, the overall prevalence of hypertension was estimated to be 17.9% for the whole country (19.9% in urban and 15.9% in rural areas) among the population aged 25 years and above.<sup>18</sup> This compares well with the estimate derived from our meta-analysis. With the conduct of such large nationally representative surveys, the role and relevance of meta-analyses of this nature would diminish.

This meta-analysis has certain limitations. The studies included were heterogeneous in terms of the age groups studied and classification of hypertension. Interestingly, the maximum number of studies was conducted among rural populations and thus we were unable to generate an estimate for urban Bangladesh. With increasing urbanization, this segment of the population is of great relevance to NCD prevention and control.

Our meta-analysis demonstrated that 13.5% of the adult rural population suffers from hypertension and is at risk for developing cardiovascular and kidney diseases. The study thus underscores the importance of screening asymptomatic individuals. Every adult's blood pressure should be checked. Primary health-care centres, community clinics and all health-care facilities should implement this strategy for the prevention and control of hypertension.

## References

- (1) Lim SS, Vos T, Flaxman AD, Danaei G, Shibuya K, Adair-Rohani H, et al. A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet*. 2012; 380(9859):2224–2260.
- (2) World Health Organization. Cardiovascular diseases in developing countries. *World Health Stat Q*. 1993;46:90–150.
- (3) Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo JL Jr, et al. The seventh report of the joint national committee on prevention, detection, evaluation, and treatment of high blood pressure: the JNC 7 report. *JAMA*. 2003;289(19): 2560–2572.
- (4) Ministry of Health and Family Welfare. *Strategic plan for surveillance and prevention of non-communicable diseases in Bangladesh 2007–2010*. Dhaka: Director General of Health Service, 2007.
- (5) Zaman MM, Rouf MA. Prevalence of hypertension in a Bangladeshi adult population. *J Hum Hypertens*. 1999;13(8):547–549.
- (6) Abu Sayeed M, Banu A, Khan AR, Hussain MZ. Prevalence of diabetes and hypertension in a rural population of Bangladesh. *Diabetes Care*. 1995;18(4):555–558.
- (7) Sayeed MA, Banu A, Malek MA, Khan AK. Blood pressure and coronary heart disease in NIDDM subjects at diagnosis: prevalence and risks in a Bangladeshi population. *Diabetes Res Clin Pr*. 1998;39(2):147–155.
- (8) Rahman M, Tondel M, Ahmad SA, Chowdhury IA, Faruquee MH, Axelson O. Hypertension and arsenic exposure in Bangladesh. *Hypertension*. 1999;33(1):74–78.
- (9) Zaman MM, Yoshiike N, Rouf MA, Syeed MH, Khan MR, Haque S, et al. Cardiovascular risk factors: distribution and prevalence in a rural population of Bangladesh. *J Cardiovasc Risk*. 2001;8(2):103–108.
- (10) Hypertension Study Group. Prevalence, awareness, treatment and control of hypertension among the elderly in Bangladesh and India: a multicentre study. *Bull World Health Organ*. 2001;79(6):490–500.

- (11) Sayeed MA, Banu A, Haq JA, Khanam PA, Mahtab H, Azad Khan AK. Prevalence of hypertension in Bangladesh: effect of socioeconomic risk factor on difference between rural and urban community. *Bangladesh Med Res Counc Bull.* 2002;28(1):7–18.
- (12) Zaman MM, Choudhury SR, Ahmed J, Numan SM, Islam MS, Yoshiike N. Non-biochemical risk factors for cardiovascular disease in general clinic-based rural population of Bangladesh. *J Epidemiol.* 2004;14(2):63–68.
- (13) Sayeed MA, Mahtab H, Khanam PA, Begum R, Banu A, Azad Khan AK. Diabetes and hypertension in pregnancy in a rural community of Bangladesh: a population-based study. *Diabetes Med.* 2005;22(9):1267–1271.
- (14) Chen Y, Factor-Litvak P, Howe GR, Parvez F, Ahsan H. Nutritional influence on risk of high blood pressure in Bangladesh: a population-based cross-sectional study. *Am J Clin Nutr.* 2006;84(5):1224–1232.
- (15) Zaman MM, Ahmed J, Choudhury SR, Numan SM, Parvin K, Islam MS. Prevalence of ischemic heart disease in a rural population of Bangladesh. *Indian Heart J.* 2007;59(3):239–241.
- (16) Van Minh H, Soonthornthada K, Ng N, Juvekar S, Razzaque A, Ashraf A, et al. Blood pressure in adult rural INDEPTH population in Asia. *Glob Health Action.* 2009;2.
- (17) Chen Y, Factor-Litvak P, Howe GR, Graziano JH, Brandt-Rauf P, Parvez F, et al. Arsenic exposure from drinking water, dietary intakes of B vitamins and folate, and risk of high blood pressure in Bangladesh: a population-based, cross-sectional study. *Am J Epidemiol.* 2007;165(5):541–552.
- (18) World Health Organization, Ministry of Health and Family Welfare, Bangladesh. *Non-communicable disease risk factor survey Bangladesh 2010.* Geneva, 2011.

